

Nearly Tri-bimaximal Neutrino Mixing and CP Violation from μ - τ Symmetry Breaking

Zhi-zhong Xing^{*}, He Zhang[†], Shun Zhou[‡]

*CCAST (World Laboratory), P.O. Box 8730, Beijing 100080, China
and Institute of High Energy Physics, Chinese Academy of Sciences,
P.O. Box 918, Beijing 100049, China*

Abstract

Assuming the Majorana nature of massive neutrinos, we generalize the Friedberg-Lee neutrino mass model to include CP violation in the neutrino mass matrix M_ν . We show that a favorable neutrino mixing pattern (with $\theta_{12} \approx 35.3^\circ$, $\theta_{23} = 45^\circ$, $\theta_{13} \neq 0^\circ$ and $\delta = 90^\circ$) can naturally be derived from M_ν , if it has an approximate or softly-broken μ - τ symmetry. We point out a different way to obtain the nearly tri-bimaximal neutrino mixing with $\delta = 0^\circ$ and non-vanishing Majorana phases. The most general case, in which all the free parameters of M_ν are complex and the resultant neutrino mixing matrix contains both Dirac and Majorana phases of CP violation, is also discussed.

PACS number(s): 14.60.Lm, 14.60.Pq, 95.85.Ry

Typeset using REVTeX

^{*}E-mail: xingzz@mail.ihep.ac.cn

[†]E-mail: zhanghe@mail.ihep.ac.cn

[‡]E-mail: zhoush@mail.ihep.ac.cn

I. INTRODUCTION

The solar [1], atmospheric [2], reactor [3] and accelerator [4] neutrino experiments have provided us with very convincing evidence that neutrinos are massive and lepton flavors are mixed. Given the basis in which the flavor eigenstates of charged leptons are identified with their mass eigenstates, the phenomenon of neutrino mixing can simply be described by a 3×3 unitary matrix V which transforms the neutrino mass eigenstates (ν_1, ν_2, ν_3) into the neutrino flavor eigenstates $(\nu_e, \nu_\mu, \nu_\tau)$. We assume massive neutrinos to be Majorana particles and parametrize V as

$$V = \begin{pmatrix} c_{12}c_{13} & s_{12}c_{13} & s_{13}e^{-i\delta} \\ -s_{12}c_{23} - c_{12}s_{23}s_{13}e^{i\delta} & c_{12}c_{23} - s_{12}s_{23}s_{13}e^{i\delta} & s_{23}c_{13} \\ s_{12}s_{23} - c_{12}c_{23}s_{13}e^{i\delta} & -c_{12}s_{23} - s_{12}c_{23}s_{13}e^{i\delta} & c_{23}c_{13} \end{pmatrix} \begin{pmatrix} e^{i\rho} & 0 & 0 \\ 0 & e^{i\sigma} & 0 \\ 0 & 0 & 1 \end{pmatrix}, \quad (1)$$

where $c_{ij} \equiv \cos \theta_{ij}$ and $s_{ij} \equiv \sin \theta_{ij}$ (for $ij = 12, 23$ and 13). A global analysis of current experimental data [5] yields $30^\circ < \theta_{12} < 38^\circ$, $36^\circ < \theta_{23} < 54^\circ$ and $\theta_{13} < 10^\circ$ at the 99% confidence level, but three CP-violating phases of V (i.e., the Dirac phase δ and the Majorana phases ρ and σ) are entirely unrestricted. In order to interpret the largeness of two neutrino mixing angles together with the smallness of three neutrino masses, many theoretical and phenomenological models of lepton mass matrices have been proposed in the literature [6]. Among them, the scenarios based on possible flavor symmetries are particularly simple, suggestive and predictive.

In this paper, we focus our interest on the neutrino mass model proposed recently by Friedberg and Lee (FL) [7]. The neutrino mass operator in the FL model is simply given by

$$\begin{aligned} \mathcal{L}_{\nu\text{-mass}} = & a (\bar{\nu}_\tau - \bar{\nu}_\mu) (\nu_\tau - \nu_\mu) + b (\bar{\nu}_\mu - \bar{\nu}_e) (\nu_\mu - \nu_e) + c (\bar{\nu}_e - \bar{\nu}_\tau) (\nu_e - \nu_\tau) \\ & + m_0 (\bar{\nu}_e \nu_e + \bar{\nu}_\mu \nu_\mu + \bar{\nu}_\tau \nu_\tau), \end{aligned} \quad (2)$$

where a, b, c and m_0 are all assumed to be *real*. A salient feature of $\mathcal{L}_{\nu\text{-mass}}$ is its partial gauge-like symmetry; i.e., its a, b and c terms are invariant under the transformation $\nu_\alpha \rightarrow \nu_\alpha + z$ (for $\alpha = e, \mu, \tau$) with z being a space-time independent constant element of the Grassmann algebra [7]. Eq. (2) means that the neutrino mass matrix M_ν takes the form

$$M_\nu = m_0 \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} + \begin{pmatrix} b+c & -b & -c \\ -b & a+b & -a \\ -c & -a & a+c \end{pmatrix}. \quad (3)$$

Diagonalizing M_ν by the transformation $V_{\text{FL}}^\dagger M_\nu V_{\text{FL}}^* = \text{Diag}\{m_1, m_2, m_3\}$, in which m_i (for $i = 1, 2, 3$) stand for the neutrino masses, one may obtain the neutrino mixing matrix

$$V_{\text{FL}} = \begin{pmatrix} \frac{2}{\sqrt{6}} & \frac{1}{\sqrt{3}} & 0 \\ -\frac{1}{\sqrt{6}} & \frac{1}{\sqrt{3}} & \frac{1}{\sqrt{2}} \\ -\frac{1}{\sqrt{6}} & \frac{1}{\sqrt{3}} & -\frac{1}{\sqrt{2}} \end{pmatrix} \begin{pmatrix} \cos \frac{\theta}{2} & 0 & \sin \frac{\theta}{2} \\ 0 & 1 & 0 \\ -\sin \frac{\theta}{2} & 0 & \cos \frac{\theta}{2} \end{pmatrix}, \quad (4)$$

where θ is given by $\tan \theta = \sqrt{3}(b-c)/[(b+c)-2a]$. This interesting result leads us to the following observations:

- If $\theta = 0^\circ$ holds, V_{FL} will reproduce the exact tri-bimaximal neutrino mixing pattern (with $\tan \theta_{12} = 1/\sqrt{2}$ or $\theta_{12} \approx 35.3^\circ$, $\theta_{23} = 45^\circ$ and $\theta_{13} = 0^\circ$) [8]. The latter, which can be understood as a geometric representation of the neutrino mixing matrix [9], is in good agreement with current experimental data. Non-vanishing but small θ predicts $\sin \theta_{13} = (2/\sqrt{6}) \sin(\theta/2)$, implying $\theta \lesssim 24.6^\circ$ for $\theta_{13} < 10^\circ$. On the other hand, θ_{23} will mildly deviate from its best-fit value $\theta_{23} = 45^\circ$ if θ (or θ_{13}) takes non-zero values.
- The limit $\theta = 0^\circ$ results from $b = c$. When $b = c$ holds, it is easy to check that the neutrino mass operator $\mathcal{L}_{\nu-\text{mass}}$ has the exact μ - τ symmetry (i.e., $\mathcal{L}_{\nu-\text{mass}}$ is invariant under the exchange of μ and τ indices). In other words, the tri-bimaximal neutrino mixing is a natural consequence of the μ - τ symmetry of M_ν in the FL model. Then $\theta_{13} \neq 0^\circ$ and $\theta_{23} \neq 45^\circ$ measure the strength of μ - τ symmetry breaking, as many authors have discussed in other neutrino mass models [10].

In addition, one may consider to remove one degree of freedom from $\mathcal{L}_{\nu-\text{mass}}$ or M_ν (for instance, by setting $c = 0$ [7]).

We aim to generalize the FL model to accommodate CP and T violation for massive Majorana neutrinos ¹. The effective Majorana neutrino mass term can be written as

$$\mathcal{L}'_{\nu-\text{mass}} = \frac{1}{2} \overline{(\nu_e, \nu_\mu, \nu_\tau)_L} M_\nu \begin{pmatrix} \nu_e^c \\ \nu_\mu^c \\ \nu_\tau^c \end{pmatrix}_R + \text{h.c.} , \quad (5)$$

where $\nu_\alpha^c \equiv C\bar{\nu}_\alpha^T$ (for $\alpha = e, \mu, \tau$), and M_ν is of the same form as that given in Eq. (3). Now the parameters of M_ν (i.e., a, b, c and m_0) are all complex. Then we are able to derive both the Dirac phase (δ) and the Majorana phases (ρ and σ) for the neutrino mixing matrix V . Two special cases are particularly interesting:

- Scenario (A): a and m_0 are real, and $b = c^*$ are complex. We find that the μ - τ symmetry of M_ν is softly broken in this case, leading to the elegant predictions $\theta_{13} \neq 0^\circ$, $\theta_{23} = 45^\circ$ and $\delta = 90^\circ$. Two Majorana phases ρ and σ keep vanishing.
- Scenario (B): a, b and c are real, but m_0 is complex. We find that the results of θ_{12} , θ_{23} and θ_{13} obtained from V_{FL} keep unchanged in this case, but some nontrivial values of the Majorana phases ρ and σ can now be generated. The Dirac phase δ remains vanishing.

The remaining part of this paper is organized as follows. More detailed discussions about scenarios (A) and (B) will be presented in section II. Section III is devoted to a generic analysis of the generalized FL model with no special assumptions. Finally, we make some concluding remarks in section IV.

¹To include T violation into the model, Friedberg and Lee [7] have inserted the phase factors $e^{\pm i\eta}$ into Eq. (2) by replacing the term $c(\bar{\nu}_e - \bar{\nu}_\tau)(\nu_e - \nu_\tau)$ with the term $c(e^{-i\eta}\bar{\nu}_e - \bar{\nu}_\tau)(e^{+i\eta}\nu_e - \nu_\tau)$. The resultant neutrino mass matrix is no longer symmetric, hence it definitely describes Dirac neutrinos instead of Majorana neutrinos.

II. TWO SIMPLE SCENARIOS

First of all, let us consider two special but interesting scenarios of the generalized FL model and explore their respective consequences on three neutrino mixing angles and three CP-violating phases. They are quite instructive in phenomenology and may easily be tested by a variety of long-baseline neutrino oscillation experiments in the near future.

A. Scenario (A)

In this scenario, a and m_0 are real, b and c are complex, and $b = c^*$ holds. The corresponding neutrino mass matrix M_ν reads

$$M_\nu = m_0 \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} + \begin{pmatrix} 2\text{Re}(b) & -b & -b^* \\ -b & a+b & -a \\ -b^* & -a & a+b^* \end{pmatrix}. \quad (6)$$

We find that M_ν can be diagonalized by the transformation $V^\dagger M_\nu V^* = \text{Diag}\{m_1, m_2, m_3\}$, where

$$V = \begin{pmatrix} \frac{2}{\sqrt{6}} & \frac{1}{\sqrt{3}} & 0 \\ -\frac{1}{\sqrt{6}} & \frac{1}{\sqrt{3}} & \frac{1}{\sqrt{2}} \\ -\frac{1}{\sqrt{6}} & \frac{1}{\sqrt{3}} & -\frac{1}{\sqrt{2}} \end{pmatrix} \begin{pmatrix} \cos \frac{\theta}{2} & 0 & \sin \frac{\theta}{2} e^{-i\delta} \\ 0 & 1 & 0 \\ -\sin \frac{\theta}{2} e^{i\delta} & 0 & \cos \frac{\theta}{2} \end{pmatrix}, \quad (7)$$

and δ is just the Dirac phase of CP violation defined in the standard parametrization of V . A straightforward calculation yields $\delta = 90^\circ$,

$$\tan \theta = \frac{\sqrt{3} \text{Im}(b)}{m_0 + a + 2\text{Re}(b)}, \quad (8)$$

together with three neutrino masses

$$\begin{aligned} m_1 &= \sqrt{[m_0 + a + 2\text{Re}(b)]^2 + 3[\text{Im}(b)]^2} - a + \text{Re}(b), \\ m_2 &= m_0, \\ m_3 &= \sqrt{[m_0 + a + 2\text{Re}(b)]^2 + 3[\text{Im}(b)]^2} + a - \text{Re}(b). \end{aligned} \quad (9)$$

Note that the special result of δ is a natural consequence of the purely imaginary term $b - c = 2i\text{Im}(b)$ in this scenario. The difference between b and c can be referred to as the *soft* μ - τ symmetry breaking, because $|b| = |c|$ holds. The explicit expression of V is

$$V = \begin{pmatrix} \frac{2}{\sqrt{6}} \cos \frac{\theta}{2} & \frac{1}{\sqrt{3}} & -i \frac{2}{\sqrt{6}} \sin \frac{\theta}{2} \\ -\frac{1}{\sqrt{6}} \cos \frac{\theta}{2} - i \frac{1}{\sqrt{2}} \sin \frac{\theta}{2} & \frac{1}{\sqrt{3}} & \frac{1}{\sqrt{2}} \cos \frac{\theta}{2} + i \frac{1}{\sqrt{6}} \sin \frac{\theta}{2} \\ -\frac{1}{\sqrt{6}} \cos \frac{\theta}{2} + i \frac{1}{\sqrt{2}} \sin \frac{\theta}{2} & \frac{1}{\sqrt{3}} & -\frac{1}{\sqrt{2}} \cos \frac{\theta}{2} + i \frac{1}{\sqrt{6}} \sin \frac{\theta}{2} \end{pmatrix}. \quad (10)$$

We observe that the only difference between V in Eq. (10) and V_{FL} in Eq. (4) is the introduction of a special CP-violating phase (see also Ref. [11] for a discussion about the maximal leptonic CP violation with $\delta = 90^\circ$). This CP-violating phase, which is attributed to the soft breaking of μ - τ symmetry, can change the prediction of V_{FL} for θ_{23} . Comparing Eq. (10) with Eq. (1), we immediately obtain ²

$$\begin{aligned}\sin \theta_{12} &= \frac{1}{\sqrt{2 + \cos \theta}}, \\ \sin \theta_{23} &= \frac{1}{\sqrt{2}}, \\ \sin \theta_{13} &= \frac{2}{\sqrt{6}} \sin \frac{\theta}{2},\end{aligned}\tag{11}$$

together with $\delta = 90^\circ$ and $\rho = \sigma = 0^\circ$. Without loss of generality, we have restricted θ to the first quadrant. The leptonic Jarlskog parameter $\mathcal{J}$ [12], which is a rephasing-invariant measure of CP violation in neutrino oscillations, reads $\mathcal{J} = \sin \theta / (6\sqrt{3})$. One can see that the soft breaking of μ - τ symmetry leads to both $\theta_{13} \neq 0^\circ$ and $\mathcal{J} \neq 0$, but it does not affect the favorable result $\theta_{23} = 45^\circ$ given by the tri-bimaximal neutrino mixing pattern. On the other hand, $\sin \theta_{12} \approx 1/\sqrt{3}$ is an excellent approximation, since θ must be small to maintain the smallness of θ_{13} . In view of $\theta_{13} < 10^\circ$, we obtain $\theta \lesssim 24.6^\circ$ and $\mathcal{J} \lesssim 0.04$. It is possible to measure $\mathcal{J} \sim \mathcal{O}(10^{-2})$ in the future long-baseline neutrino oscillation experiments.

Note that the neutrino masses rely on four real model parameters m_0 , a , $\text{Re}(b)$ and $\text{Im}(b)$. Thus it is easy to fit the neutrino mass-squared differences $\Delta m_{21}^2 = (7.2 \cdots 8.9) \times 10^{-5} \text{ eV}^2$ and $\Delta m_{32}^2 = \pm(2.1 \cdots 3.1) \times 10^{-3} \text{ eV}^2$ [5]. Such a fit should not involve any fine-tuning, because (a) the number of free parameters is larger than the number of constraint conditions and (b) three neutrino masses have very weak correlation with three mixing angles. A detailed numerical analysis shows that only the normal mass hierarchy ($m_1 < m_2 < m_3$) is allowed in this scenario. FIG. 1 illustrates the parameter space of m_0 , a , $\text{Re}(b)$ and $\text{Im}(b)$, where $m_0 \lesssim 0.2 \text{ eV}$ has typically been taken as a generous upper bound on the absolute neutrino mass [13]. Because of $m_0 = m_2$, it is straightforward to get $m_0 > \sqrt{\Delta m_{21}^2} \approx 0.009 \text{ eV}$, as shown in FIG. 1. The small μ - τ symmetry breaking (or small θ) requires the small magnitude of $\text{Im}(b)$. Thus Eq. (10) allows us to get $\text{Re}(b) \approx (m_1 - m_2)/3$ in the $\text{Im}(b) \rightarrow 0$ limit, implying that $\text{Re}(b)$ is negative and its magnitude is very small. Furthermore, $a \approx (m_3 - m_2)/2$ approximately holds in the $\text{Im}(b) \rightarrow 0$ limit, implying that the maximal value of a is roughly half of $m_3 \sim \sqrt{|\Delta m_{32}^2|}$ (i.e., $a \lesssim 0.025 \text{ eV}$) when three neutrino masses develop a strong normal hierarchy.

B. Scenario (B)

In this simple scenario, only m_0 is assumed to be complex. While the neutrino mass matrix M_ν takes the same form as that given in Eq. (3), its diagonalization ($V^\dagger M_\nu V^* =$

²It is trivial to redefine the phases of charged-lepton fields, such that the location of CP-violating phases in Eq. (10) is the same as that in Eq. (1).

$\text{Diag}\{m_1, m_2, m_3\}$) requires the following transformation matrix

$$V = \begin{pmatrix} \frac{2}{\sqrt{6}} & \frac{1}{\sqrt{3}} & 0 \\ -\frac{1}{\sqrt{6}} & \frac{1}{\sqrt{3}} & \frac{1}{\sqrt{2}} \\ -\frac{1}{\sqrt{6}} & \frac{1}{\sqrt{3}} & -\frac{1}{\sqrt{2}} \end{pmatrix} \begin{pmatrix} \cos \frac{\theta}{2} & 0 & \sin \frac{\theta}{2} \\ 0 & 1 & 0 \\ -\sin \frac{\theta}{2} & 0 & \cos \frac{\theta}{2} \end{pmatrix} \begin{pmatrix} e^{i\phi_1} & 0 & 0 \\ 0 & e^{i\phi_2} & 0 \\ 0 & 0 & e^{i\phi_3} \end{pmatrix}, \quad (12)$$

where ϕ_i (for $i = 1, 2, 3$) originate from the imaginary part of m_0 . Comparing between Eqs. (1) and (12), one can see that the Majorana phases of CP violation in the standard parametrization of V are given by $\rho = \phi_1 - \phi_3$ and $\sigma = \phi_2 - \phi_3$ (i.e., the phase factor $e^{i\phi_3}$ in Eq. (12) is finally rotated away by redefining the phases of three charged-lepton fields). After a straightforward calculation, we obtain

$$\tan \theta = \frac{\sqrt{3}(b-c)}{(b+c)-2a}, \quad (13)$$

and

$$\begin{aligned} m_1 &= \sqrt{[m_- + \text{Re}(m_0)]^2 + [\text{Im}(m_0)]^2}, \\ m_2 &= |m_0|, \\ m_3 &= \sqrt{[m_+ + \text{Re}(m_0)]^2 + [\text{Im}(m_0)]^2}, \end{aligned} \quad (14)$$

where

$$m_{\pm} = (a+b+c) \pm \sqrt{a^2 + b^2 + c^2 - ab - ac - bc}. \quad (15)$$

The final result for the neutrino mixing matrix V is

$$V = \begin{pmatrix} \frac{2}{\sqrt{6}} \cos \frac{\theta}{2} & \frac{1}{\sqrt{3}} & \frac{2}{\sqrt{6}} \sin \frac{\theta}{2} \\ -\frac{1}{\sqrt{6}} \cos \frac{\theta}{2} - \frac{1}{\sqrt{2}} \sin \frac{\theta}{2} & \frac{1}{\sqrt{3}} & \frac{1}{\sqrt{2}} \cos \frac{\theta}{2} - \frac{1}{\sqrt{6}} \sin \frac{\theta}{2} \\ -\frac{1}{\sqrt{6}} \cos \frac{\theta}{2} + \frac{1}{\sqrt{2}} \sin \frac{\theta}{2} & \frac{1}{\sqrt{3}} & -\frac{1}{\sqrt{2}} \cos \frac{\theta}{2} - \frac{1}{\sqrt{6}} \sin \frac{\theta}{2} \end{pmatrix} \begin{pmatrix} e^{i\rho} & 0 & 0 \\ 0 & e^{i\sigma} & 0 \\ 0 & 0 & 1 \end{pmatrix}, \quad (16)$$

where ρ and σ are given by

$$\begin{aligned} \tan 2\rho &= \frac{(m_+ - m_-) \text{Im}(m_0)}{|m_0|^2 + m_+ m_- + (m_+ + m_-) \text{Re}(m_0)}, \\ \tan 2\sigma &= \frac{m_+ \text{Im}(m_0)}{|m_0|^2 + m_+ \text{Re}(m_0)}. \end{aligned} \quad (17)$$

One can see that the only difference between V in Eq. (16) and V_{FL} in Eq. (4) is the introduction of two Majorana phases of CP violation. Although ρ and σ have nothing to

do with the behaviors of neutrino oscillations, they may significantly affect the neutrinoless double-beta decay [14]. Comparing Eq. (16) with Eq. (1), we arrive at

$$\begin{aligned}\sin \theta_{12} &= \frac{1}{\sqrt{2 + \cos \theta}}, \\ \sin \theta_{23} &= \frac{\sqrt{2 + \cos \theta - \sqrt{3} \sin \theta}}{\sqrt{2(2 + \cos \theta)}}, \\ \sin \theta_{13} &= \frac{2}{\sqrt{6}} \sin \frac{\theta}{2},\end{aligned}\tag{18}$$

together with $\delta = 0^\circ$ for the Dirac phase of CP violation. Again θ has been restricted to the first quadrant. The results for θ_{12} and θ_{13} in this scenario are the same as those obtained in scenario (A), but the Jarlskog parameter $\mathcal{J}$ is now vanishing. Because of the μ - τ symmetry breaking, θ_{23} may somehow deviate from the favorable value $\theta_{23} = 45^\circ$. Given $\theta \lesssim 24.6^\circ$ corresponding to $\theta_{13} < 10^\circ$, θ_{23} is allowed to vary in the range $37.8^\circ \lesssim \theta_{23} \leq 45^\circ$ [15].

Note that the neutrino masses depend on five real model parameters a , b , c , $\text{Re}(m_0)$ and $\text{Im}(m_0)$. Hence there is sufficient freedom to fit two observed neutrino mass-squared differences Δm_{21}^2 and Δm_{32}^2 . A detailed numerical analysis shows that both the normal mass hierarchy ($m_1 < m_2 < m_3$) and the inverted mass hierarchy ($m_3 < m_1 < m_2$) are allowed in scenario (B). FIGs. 2 and 3 illustrate the parameter space of a , b , c , $\text{Re}(m_0)$ and $\text{Im}(m_0)$, where $|m_0| \lesssim 0.2$ eV has been taken [13]. Taking account of the analytical reciprocity between b and c (i.e., an exchange of b and c in Eq. (15) does not affect the result of m_i obtained in Eq. (14)), we plot their allowed regions in the same figure without any confusion. Note that m_0 cannot be purely imaginary in both normal and inverted mass hierarchies, otherwise we would be left with $\Delta m_{21}^2 < 0$, which is in disagreement with current experimental data. Moreover, the lower bound of $|\text{Re}(m_0)|$ is obviously restricted by the magnitude of m_2 (i.e., $m_2 \sim \sqrt{\Delta m_{21}^2} \approx 0.009$ eV in the normal mass hierarchy or $m_2 \sim \sqrt{|\Delta m_{32}^2|} \approx 0.05$ eV in the inverted mass hierarchy). In view of $(m_+ - m_-) \propto (m_3^2 - m_1^2)$ from Eq. (14), we find $m_+ \neq m_-$. This inequality implies that $a = b = c$ is not allowed, as one can see from Eq. (15) or FIGs. 2 and 3. The disconnected regions in the plots of a versus b (or c) are ascribed to the ambiguity induced by the sign of $\text{Re}(m_0)$. FIG. 4 shows the allowed regions of ρ and σ . We see that both of them are less restricted, as a consequence of the large freedom associated with the imaginary part of m_0 .

III. GENERIC ANALYSIS

Now let us assume all the parameters of M_ν in Eq. (3) to be complex and calculate its mass eigenvalues and flavor mixing parameters. Using the tri-bimaximal mixing matrix [8]

$$V_0 = \begin{pmatrix} \frac{2}{\sqrt{6}} & \frac{1}{\sqrt{3}} & 0 \\ -\frac{1}{\sqrt{6}} & \frac{1}{\sqrt{3}} & \frac{1}{\sqrt{2}} \\ -\frac{1}{\sqrt{6}} & \frac{1}{\sqrt{3}} & -\frac{1}{\sqrt{2}} \end{pmatrix},\tag{19}$$

we transform M_ν into the following form:

$$V_0^\dagger M_\nu V_0^* = \begin{pmatrix} \frac{3}{2}(b+c) + m_0 & 0 & \frac{\sqrt{3}}{2}(c-b) \\ 0 & m_0 & 0 \\ \frac{\sqrt{3}}{2}(c-b) & 0 & 2a + \frac{1}{2}(b+c) + m_0 \end{pmatrix}. \quad (20)$$

This complex mass matrix can be diagonalized by the transformation $V_1^\dagger (V_0^\dagger M_\nu V_0^*) V_1^* = \text{Diag}\{m_1, m_2, m_3\}$, where

$$V_1 = \begin{pmatrix} \cos \frac{\theta}{2} & 0 & \sin \frac{\theta}{2} e^{-i\delta} \\ 0 & 1 & 0 \\ -\sin \frac{\theta}{2} e^{i\delta} & 0 & \cos \frac{\theta}{2} \end{pmatrix} \begin{pmatrix} e^{i\phi_1} & 0 & 0 \\ 0 & e^{i\phi_2} & 0 \\ 0 & 0 & e^{i\phi_3} \end{pmatrix}. \quad (21)$$

The neutrino mixing matrix V turns out to be $V = V_0 V_1$, in which δ is just the Dirac phase of CP violation. Of course, $\rho = \phi_1 - \phi_3$ and $\sigma = \phi_2 - \phi_3$ are the Majorana phases of CP violation in the standard parametrization of V .

To be explicit, we express θ and δ in terms of the parameters of M_ν . The results are

$$\begin{aligned} \tan \theta &= \frac{\sqrt{X^2 + Y^2}}{Z}, \\ \tan \delta &= \frac{X}{Y}, \end{aligned} \quad (22)$$

where

$$\begin{aligned} X &= \sqrt{3} \text{Im}(ac^* - ab^* - bc^*), \\ Y &= \sqrt{3} [|c|^2 - |b|^2 + \text{Re}(ac^* - ab^* - m_0 b^* + m_0 c^*)], \\ Z &= 2|a|^2 - |b|^2 - |c|^2 + \text{Re}(ab^* + ac^* - 2bc^* + 2m_0 a^* - m_0 b^* - m_0 c^*). \end{aligned} \quad (23)$$

Furthermore, three mass eigenvalues of M_ν and two Majorana phases of V are found to be

$$\begin{aligned} m_1 &= \left| T_1 \cos^2 \frac{\theta}{2} - T_2 \sin \theta e^{-i\delta} + T_3 \sin^2 \frac{\theta}{2} e^{-i2\delta} \right|, \\ m_2 &= |m_0|, \\ m_3 &= \left| T_3 \cos^2 \frac{\theta}{2} + T_2 \sin \theta e^{+i\delta} + T_1 \sin^2 \frac{\theta}{2} e^{+i2\delta} \right|; \end{aligned} \quad (24)$$

and

$$\begin{aligned} \rho &= \frac{1}{2} \arg \left(\frac{T_1 \cos^2 \frac{\theta}{2} - T_2 \sin \theta e^{-i\delta} + T_3 \sin^2 \frac{\theta}{2} e^{-i2\delta}}{T_3 \cos^2 \frac{\theta}{2} + T_2 \sin \theta e^{+i\delta} + T_1 \sin^2 \frac{\theta}{2} e^{+i2\delta}} \right), \\ \sigma &= \frac{1}{2} \arg \left(\frac{m_0}{T_3 \cos^2 \frac{\theta}{2} + T_2 \sin \theta e^{+i\delta} + T_1 \sin^2 \frac{\theta}{2} e^{+i2\delta}} \right), \end{aligned} \quad (25)$$

where

$$\begin{aligned}
T_1 &= \frac{3}{2}(b+c) + m_0, \\
T_2 &= \frac{\sqrt{3}}{2}(c-b), \\
T_3 &= 2a + \frac{1}{2}(b+c) + m_0.
\end{aligned} \tag{26}$$

The final result of V is

$$V = \begin{pmatrix} \frac{2}{\sqrt{6}} \cos \frac{\theta}{2} & \frac{1}{\sqrt{3}} & \frac{2}{\sqrt{6}} \sin \frac{\theta}{2} e^{-i\delta} \\ -\frac{1}{\sqrt{6}} \cos \frac{\theta}{2} - \frac{1}{\sqrt{2}} \sin \frac{\theta}{2} e^{i\delta} & \frac{1}{\sqrt{3}} & \frac{1}{\sqrt{2}} \cos \frac{\theta}{2} - \frac{1}{\sqrt{6}} \sin \frac{\theta}{2} e^{-i\delta} \\ -\frac{1}{\sqrt{6}} \cos \frac{\theta}{2} + \frac{1}{\sqrt{2}} \sin \frac{\theta}{2} e^{i\delta} & \frac{1}{\sqrt{3}} & -\frac{1}{\sqrt{2}} \cos \frac{\theta}{2} - \frac{1}{\sqrt{6}} \sin \frac{\theta}{2} e^{-i\delta} \end{pmatrix} \begin{pmatrix} e^{i\rho} & 0 & 0 \\ 0 & e^{i\sigma} & 0 \\ 0 & 0 & 1 \end{pmatrix}, \tag{27}$$

from which we obtain the Jarlskog parameter $\mathcal{J} = \sin \theta \sin \delta / (6\sqrt{3})$. Comparing Eq. (27) with Eq. (1), we arrive at

$$\begin{aligned}
\sin \theta_{12} &= \frac{1}{\sqrt{2 + \cos \theta}}, \\
\sin \theta_{23} &= \frac{\sqrt{2 + \cos \theta - \sqrt{3} \sin \theta \cos \delta}}{\sqrt{2(2 + \cos \theta)}}, \\
\sin \theta_{13} &= \frac{2}{\sqrt{6}} \sin \frac{\theta}{2}.
\end{aligned} \tag{28}$$

One can see that the predictions for θ_{12} and θ_{13} are the same as those obtained in scenarios (A) and (B). Thus they are two typical and general consequences of the FL model. As for θ_{23} , the generic result in Eq. (28) may easily reproduce the special result in Eq. (11) for scenario (A) with $\delta = 90^\circ$ or that in Eq. (18) for scenario (B) with $\delta = 0^\circ$. Given $\theta \lesssim 24.6^\circ$ (i.e., $\theta_{13} < 10^\circ$), θ_{23} is allowed to vary in the range $37.8^\circ \leq \theta_{23} \lesssim 52.2^\circ$ for arbitrary δ .

As m_0 , a , b and c are all complex, we now have much more freedom to fit two observed neutrino mass-squared differences Δm_{21}^2 and Δm_{32}^2 . Of course, both the normal neutrino mass hierarchy ($m_1 < m_2 < m_3$) and the inverted neutrino mass hierarchy ($m_3 < m_1 < m_2$) are expected in this generic case. We shall not carry out a numerical analysis of the parameter space, however, just because it involves a lot of uncertainties and is not as suggestive as that in scenario (A) or scenario (B).

IV. CONCLUDING REMARKS

We have pointed out that the nearly tri-bimaximal neutrino mixing can be regarded as a natural consequence of the slight μ - τ symmetry breaking in the FL neutrino mass model. Another straightforward consequence of the μ - τ symmetry breaking is CP violation.

Assuming the Majorana nature of massive neutrinos, we have generalized the FL model to introduce the CP-violating effects. In addition to a generic analysis of the generalized FL model, two simple but intriguing scenarios have been proposed: scenario (A) involves the softly-broken μ - τ symmetry, leading to the elegant predictions $\theta_{13} \neq 0^\circ$, $\theta_{23} = 45^\circ$, $\delta = 90^\circ$ and vanishing Majorana phases of CP violation; scenario (B) predicts $\theta_{13} \neq 0^\circ$, $\theta_{23} \neq 45^\circ$, $\delta = 0^\circ$ and two nontrivial Majorana phases of CP violation, on the other hand. Both scenarios are compatible with current experimental data.

Although our discussions about the generalized FL model are restricted to low-energy scales, it can certainly be extended to a superhigh-energy scale (e.g., the GUT scale or the seesaw scale). In this case, one should take into account the radiative corrections to both neutrino masses and flavor mixing parameters when they run from the high scale to the electroweak scale. A particularly interesting point is that the Majorana phases ρ and σ can radiatively be generated from the Dirac phase δ in scenario (A), while the Dirac phase δ can radiatively be generated from the Majorana phases ρ and σ in scenario (B) [16]. Thus the features of leptonic CP violation can fully show up in both scenarios at low-energy scales, if they are originally prescribed at a superhigh-energy scale (see Ref. [17] for a detailed analysis of the running behaviors of CP-violating phases in the nearly tri-bimaximal neutrino mixing pattern, including the minimal supergravity threshold effects).

We conclude that the μ - τ symmetry and its slight breaking are useful and suggestive for model building. We expect that a stringent test of the generalized FL model, in particular its two simple and instructive scenarios, can be achieved in the near future from the reactor neutrino oscillation experiments (towards measuring θ_{13}), the accelerator long-baseline neutrino oscillation experiments (towards detecting both θ_{13} and δ) and the neutrinoless double-beta decay experiments (towards probing the Majorana nature of massive neutrinos and constraining their Majorana phases of CP violation).

ACKNOWLEDGMENTS

We are indebted to W.Q. Chao for bringing Refs. [7] and [9] to our particular attention. One of us (Z.Z.X.) would also like to thank T.D. Lee for his encouragement at the Workshop on Future China-US Cooperation in High Energy Physics (June 2006, Beijing). This work is supported in part by the National Natural Science Foundation of China.

REFERENCES

- [1] SNO Collaboration, Q.R. Ahmad *et al.*, Phys. Rev. Lett. **89**, 011301 (2002).
- [2] For a review, see: C.K. Jung *et al.*, Ann. Rev. Nucl. Part. Sci. **51**, 451 (2001).
- [3] KamLAND Collaboration, K. Eguchi *et al.*, Phys. Rev. Lett. **90**, 021802 (2003).
- [4] K2K Collaboration, M.H. Ahn *et al.*, Phys. Rev. Lett. **90**, 041801 (2003).
- [5] A. Strumia and F. Vissani, hep-ph/0606054.
- [6] For recent reviews with extensive references, see: H. Fritzsch and Z.Z. Xing, Prog. Part. Nucl. Phys. **45**, 1 (2000); Z.Z. Xing, Int. J. Mod. Phys. A **19**, 1 (2004); Altarelli and F. Feruglio, New J. Phys. **6**, 106 (2004); R.N. Mohapatra *et al.*, hep-ph/0510213; R.N. Mohapatra and A.Yu. Smirnov, hep-ph/0603118; A. Strumia and F. Vissani, hep-ph/0606054.
- [7] R. Friedberg and T.D. Lee, HEP&NP **30**, 591 (2006), hep-ph/0606071; T.D. Lee, invited talk given at the Workshop on Future China-US Cooperation in High Energy Physics, Beijing, June 11–18, 2006.
- [8] P.F. Harrison, D.H. Perkins, and W.G. Scott, Phys. Lett. B **530**, 167 (2002); Z.Z. Xing, Phys. Lett. B **533**, 85 (2002); P.F. Harrison and W.G. Scott, Phys. Lett. B **535**, 163 (2002); X.G. He and A. Zee, Phys. Lett. B **560**, 87 (2003); C.I. Low and R.R. Volkas, Phys. Rev. D **68**, 033007 (2003); E. Ma, Phys. Lett. B **583**, 157 (2004); hep-ph/0409075; G. Altarelli and F. Feruglio, Nucl. Phys. B **720**, 64 (2005); F. Plentinger and W. Rodejohann, Phys. Lett. B **625**, 264 (2005); K.S. Babu and X.G. He, hep-ph/0507217; A. Zee, Phys. Lett. B **630**, 58 (2005); S.K. Kang, Z.Z. Xing and S. Zhou, Phys. Rev. D **73**, 013001 (2006); X.G. He, Y.Y. Keum and R.R. Volkas, JHEP **0604**, 039 (2006).
- [9] T.D. Lee, Chinese Phys. **15**, 1009 (2006), hep-ph/0605017.
- [10] See, e.g., T. Fukuyama and H. Nishiura, hep-ph/9702253; R.N. Mohapatra and S. Nussinov, Phys. Rev. D **60**, 013002 (1999); Z.Z. Xing, Phys. Rev. D **61**, 057301 (2000); Phys. Rev. D **64**, 093013 (2001); E. Ma and M. Raidal, Phys. Rev. Lett. **87**, 011802 (2001); C.S. Lam, Phys. Lett. B **507**, 214 (2001); T. Kitabayashi and M. Yasuè, Phys. Rev. D **67**, 015006 (2003); W. Grimus and L. Lavoura, Phys. Lett. B **572**, 189 (2003); J. Phys. G **30**, 73 (2004); Y. Koide, Phys. Rev. D **69**, 093001 (2004); R.N. Mohapatra, JHEP **0410**, 027 (2004); A. de Gouvea, Phys. Rev. D **69**, 093007 (2004); A. Ghosal, Mod. Phys. Lett. A **19**, 2579 (2004); W. Grimus, A.S. Joshipura, S. Kaneko, L. Lavoura, H. Sawanaka, and M. Tanimoto, Nucl. Phys. B **713**, 151 (2005); R.N. Mohapatra and W. Rodejohann, Phys. Rev. D **72**, 053001 (2005); T. Kitabayashi and M. Yasuè, Phys. Lett. B **621**, 133 (2005); R.N. Mohapatra, S. Nasri, and H.B. Yu, Phys. Lett. B **615**, 231 (2005); A.S. Joshipura, hep-ph/0512252; K. Matsuda and H. Nishiura, Phys. Rev. D **73**, 013008 (2006); R.N. Mohapatra, S. Nasri, and H.B. Yu, Phys. Lett. B **636**, 114 (2006); hep-ph/0605020; Z.Z. Xing, Phys. Rev. D **74**, 013010 (2006); K. Fuki and M. Yasue, hep-ph/0608042.
- [11] I. Aizawa, T. Kitabayashi, and M. Yasuè, Phys. Rev. D **72**, 055014 (2005); Nucl. Phys. B **728**, 220 (2005).
- [12] C. Jarlskog, Phys. Rev. Lett. **55**, 1039 (1985); D.D. Wu, Phys. Rev. D **33**, 860 (1986).
- [13] See, e.g., D.N. Spergel, astro-ph/0603449, in which the combination of WMAP and other astronomical data yields a constraint $\sum m_i < 0.68$ eV (95% C.L.).

- [14] Z.Z. Xing, Phys. Rev. D **64**, 093013 (2001); Phys. Rev. D **65**, 077302 (2002).
- [15] Note that one may also get $45^\circ \leq \theta_{23} \lesssim 52.2^\circ$, if θ takes negative values from the third or fourth quadrant. Therefore, θ_{23} is in general allowed to vary in the range $37.8^\circ \lesssim \theta_{23} \lesssim 52.2^\circ$ for $|\theta| \lesssim 24.6^\circ$. See the discussions below Eq. (28) in section III.
- [16] J.A. Casas, J.R. Espinosa, A. Ibarra, and I. Navarro, Nucl. Phys. B **573**, 652 (2000); S. Antusch, J. Kersten, M. Lindner, and M. Ratz, Nucl. Phys. B **674**, 401 (2003); S. Luo, J.W. Mei, and Z.Z. Xing, Phys. Rev. D **72**, 053014 (2005);
- [17] S. Luo and Z.Z. Xing, Phys. Lett. B **632**, 341 (2006); Phys. Lett. B **637**, 279 (2006); M. Hirsch, E. Ma, J.C. Romao, J.W.F. Valle, and A.V. del Moral, hep-ph/0606082.

FIGURES




FIG. 1. The parameter space of (m_0, a) and $(\text{Re}(b), \text{Im}(b))$ in **Scenario (A)**, where only the normal neutrino mass hierarchy is allowed.




FIG. 2. The parameter space of $(\text{Re}(m_0), \text{Im}(m_0))$ and $(a, b \text{ or } c)$ in **Scenario (B)** with the *normal* neutrino mass hierarchy.




FIG. 3. The parameter space of $(\text{Re}(m_0), \text{Im}(m_0))$ and $(a, b \text{ or } c)$ in **Scenario (B)** with the *inverted* neutrino mass hierarchy.




FIG. 4. The parameter space of ρ and σ in **Scenario (B)** with the *normal* or *inverted* neutrino mass hierarchy. Note that the same parameter space can be obtained when $\rho \rightarrow \rho \pm n\pi/2$ and $\sigma \rightarrow \sigma \pm n\pi/2$ (for $n = 1, 2, 3, \dots$), as a straightforward consequence of Eq. (17).